

## Pfizer PFE Pharmaceuticals Israel Ltd..

P.O.Box 12133 9 Shenkar St.

Herzliya Pituach, Israel 46725

Tel: 972-9-9700500 Fax: 972-9-9700501

## <u>10.10.2023: תאריך ההודעה</u>

: בעל הרישום <u>פייזר פי אף אי פרמצבטיקה ישראל בעיימ</u> מודיע על הפסקת <u>שיווק זמנית</u> של

שם התכשיר בעברית  Ketalar 50 mg/ml Injection  Solution for injection/infusion  I.M, I.V  I.M, I.V  THE CARRING (as hydrochloride)  So mg/ml  Ketamine (as hydrochloride)  So mg/ml  As the sole anaesthetic agent for diagnostic and surgical procedures. When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported. For the induction of anaesthesia prior to the administration of other general anaesthetic agents. Specific areas of application or types of procedure: When the intramuscular route of administration is preferred. Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures. Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures). Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible. Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies. Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus. Cardiac catheterization procedures. Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia annot be delaved.	110-99-21857	מספר רישום
צורת מינון  Solution for injection/infusion  I.M, I.V  Tרך מתן  Ketamine (as hydrochloride)  So mg/ml  As the sole anaesthetic agent for diagnostic and surgical procedures. When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported. For the induction of anaesthesia prior to the administration of other general anaesthetic agents. To supplement other anaesthetic agents. Specific areas of application or types of procedure: When the intramuscular route of administration is preferred. Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures. Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures). Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible. Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies. Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus. Cardiac catheterization procedures. Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm	קטאלר להזרקה 50 מ"ג/מ"ל	
אל (as hydrochloride)    Trt מתן	Ketalar 50 mg/ml Injection	שם התכשיר באנגלית
א the sole anaesthetic agent for diagnostic and surgical procedures. When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported. For the induction of anaesthesia prior to the administration of other general anaesthetic agents. To supplement other anaesthetic agents. Specific areas of application or types of procedure: When the intramuscular route of administration is preferred. Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures. Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures). Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible. Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies. Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus. Cardiac catheterization procedures. Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm	Solution for injection/infusion	צורת מינון
א the sole anaesthetic agent for diagnostic and surgical procedures.  When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported.  For the induction of anaesthesia prior to the administration of other general anaesthetic agents.  To supplement other anaesthetic agents.  Specific areas of application or types of procedure: When the intramuscular route of administration is preferred.  Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures.  Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures).  Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible.  Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies.  Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus.  Cardiac catheterization procedures.  Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm	I.M, I.V	· ·
As the sole anaesthetic agent for diagnostic and surgical procedures. When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported. For the induction of anaesthesia prior to the administration of other general anaesthetic agents. To supplement other anaesthetic agents. Specific areas of application or types of procedure: When the intramuscular route of administration is preferred. Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures. Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures). Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible. Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies. Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus. Cardiac catheterization procedures. Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm	Ketamine (as hydrochloride)	מרכיב פעיל
When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported.  For the induction of anaesthesia prior to the administration of other general anaesthetic agents.  To supplement other anaesthetic agents.  Specific areas of application or types of procedure: When the intramuscular route of administration is preferred.  Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures.  Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures).  Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible.  Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies.  Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus.  Cardiac catheterization procedures.  Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm		,
אווויזס מוומספת מיידי מיידי מיידי מיידי	As the sole anaesthetic agent for diagnostic and surgical procedures. When used by intravenous or intramuscular injection, Ketalar is best suited for short procedures. With additional doses, or by intravenous infusion, Ketalar can be used for longer procedures. If skeletal muscle relaxation is desired a muscle relaxant should be used and respiration should be supported. For the induction of anaesthesia prior to the administration of other general anaesthetic agents.  To supplement other anaesthetic agents.  Specific areas of application or types of procedure: When the intramuscular route of administration is preferred.  Debridement, painful dressings and skin grafting in burned patients as well as other superficial surgical procedures. Neurodiagnostic procedures such as pneumoencephalograms, ventriculograms, myelograms and lumbar punctures.  Diagnostic and operative procedures of the eye, ear, nose and mouth including dental extractions. (Note: eye movement may persist during ophthalmological procedures).  Anaesthesia in poor-risk patients with depression of vital functions or where depression of vital functions must be avoided if at all possible.  Orthopaedic procedures such as closed reduction, manipulation femoral pinning, amputations and biopsies.  Sigmoidoscopy and minor surgery of the anus and rectum, circumcision and pilonidal sinus.  Cardiac catheterization procedures.  Caesarian section: as an induction agent in the absence of elevated blood pressure. Anaesthesia in the asthmatic patient, either to minimise the risk of an attack of bronchospasm developing or in the presence of bronchospasm where anaesthesia cannot be delayed.	התוויה

	השיווק
21 12 22	,
31.12.23	תאריך צפי לסיום
	הפסקת השיווק
	תאריך חזרה לשיווק
	ויש למלא משבצת זאת)
	עם חזרת התכשיר
	לשיווק <b>בפועל</b> )
כן	הכללה בסל הבריאות:
	כן / לא
סיבות תפעוליות	סיבת הפסקת השיווק:
	סיבות תפעוליות /
	סיבות מסחריות
1X10 ml	גודל/י אריזה (נא
5X10 ml	לרשום את כל הגדלים
10X10 ml	הרלוונטיים להודעה
	זאת)
אין	זמינות של גודל אריזה
	אחר או חוזק אחר של
	התכשיר