



Pfizer PFE Pharmaceuticals Israel Ltd..  
P.O.Box 12133  
9 Shenkar St.  
Herzliya Pituach, Israel 46725  
Tel: 972-9-9700500 Fax: 972-9-9700501

תאריך ההודעה: 30.10.2023

בעל הרישום פייזר פי אף אי פרמצבטיקה ישראל בע"מ מודיע על הארכת הפסקת השיווק הזמנית של:

126-05-30588	מספר רישום
ליפישור 20 מ"ג	שם התכשיר בעברית
Lipitor 20 mg	שם התכשיר באנגלית
Film coated Tablets	צורת מינון
Per Os	דרך מתן
Atorvastatin (as calcium)	מרכיב פעיל
20 מ"ג	חוזק
<p>Lipitor is indicated as an adjunct to diet for reduction of elevated total cholesterol LDL- cholesterol apolipoprotein B and triglycerides and to increase HDL Cholesterol in patients with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (corresponding to types IIa and IIb of the fredrickson classification) when response to diet and other nonpharmacological measures is inadequate. Lipitor is also patients with indicated to reduce total-C and LDL -C in homozygous familial hypercholesterolemia as an adjunct to other lipid-lowering treatment (e.g. LDL apheresis) or if such treatments are unavailable.</p> <p>Pediatric patients (10-17 years of age) :</p> <p>Atorvastatin is indicated as an adjunct to diet to reduce total -C LDL-C and apo B levels in boys and postmenarchal girls 10 to 17 years of age with heterozygous familial hypercholesterolemia if adequate trial of diet therapy the following findings are after an present:</p> <p>1. LDL-C remains &gt;or = 190 mg/dl or</p> <p>2. LDL-C remains &gt; or = 160 mg/dl and: there is a positive family or two or more other history of premature cardiovascular disease CVD risk factors are present in the pediatric patient.</p> <p>Prevention of cardiovascular and/or cerebrovascular event such as MI or stroke as an adjunct to correction of other risk factors such as hypertension in patients with three or more additional risk factors or diabetes with one additional risk factor.</p> <p>In patients with clinically evident coronary heart disease Lipitor is indicated to :</p> <p>Reduce the risk of non-fatal myocardial infarction.</p> <p>Reduce the risk of fatal and non fatal stroke.</p> <p>Reduce the risk for revascularization procedures.</p> <p>Reduce the risk of hospitalization for CHF.</p> <p>Reduce the risk of angina.</p>	התוויה
13.08.2023	תאריך תחילת הפסקת השיווק

31.10.2023 15/02/2024	תאריך צפי לסיום הפסקת השיווק
	תאריך חזרה לשיווק (יש למלא משבצת זאת עם חזרת התכשיר לשיווק <b>בפועל</b> )
כן	הכללה בסל הבריאות : כן / לא
סיבות תפעוליות	סיבת הפסקת השיווק : סיבות תפעוליות / סיבות מסחריות
10, 30, 50, 100 טבליות	גודל/י אריזה (נא לרשום את כל הגדלים הרלוונטיים להודעה זאת)
Lipitor 10 mg, Lipitor 80 mg	זמינות של גודל אריזה אחר או חוזק אחר של התכשיר