

ציפרו-טבע 2מ"ג/מ"ל תמיסה לאינפוזיה**Cipro-Teva 2mg/ml**

Contains: Ciprofloxacin (as lactate) 2mg/ 1ml

עדכונים בעלון לרופא**התוויה כפי שאושרה בתעודת הרישום:****Adults:**

Broad spectrum antibiotic for infections caused by ciprofloxacin sensitive pathogens.

Children and adolescents:

- Broncho-pulmonary infections in cystic fibrosis caused by *Pseudomonas aeruginosa*
- Complicated urinary tract infections and pyelonephritis
- Inhalation anthrax (post-exposure prophylaxis and curative treatment)

Ciprofloxacin may also be used to treat severe infections in children and adolescents when there is no other alternative. Treatment should be initiated only by physicians who are experienced in the treatment of cystic fibrosis and/or severe infections in children and adolescents.

ברצוננו להודיע שהעלון לרופא עודכן. בפירוט שלהלן כלולים העדכונים העיקריים בלבד (תוספות מסומנות באדום והסרות מידע כטקסט מחוק):

4.1 Therapeutic indications

Ciprofloxacin solution for infusion is indicated for the treatment of the following infections.

Special attention should be paid to available information on resistance to ciprofloxacin before commencing therapy. Consideration should be given to official guidance on the appropriate use of antibacterial agents.

Adults

- Lower respiratory tract infections due to Gram-negative bacteria
 - exacerbations of chronic obstructive pulmonary disease. In exacerbation of chronic obstructive pulmonary disease Ciprofloxacin should be used only when it is considered inappropriate to use other antibacterial agents that are commonly recommended for the treatment of these infections.
 - broncho-pulmonary infections in cystic fibrosis or in bronchiectasis
 - pneumonia
- Chronic suppurative otitis media
- Acute exacerbation of chronic sinusitis especially if these are caused by Gram[1]negative bacteria
- Urinary tract infections
 - Acute pyelonephritis
 - Complicated pyelonephritis
 - Bacterial prostatitis
- Genital tract infections
 - Epididymo-orchitis including cases due to susceptible *Neisseria gonorrhoeae*
 - Pelvic inflammatory disease including cases due to susceptible *Neisseria gonorrhoeae*
- Infections of the gastro-intestinal tract (e.g. travellers' diarrhoea)
- Intra-abdominal infections
- Infections of the skin and soft tissue caused by Gram-negative bacteria
- Malignant external otitis
- Infections of the bones and joints
- Inhalation anthrax (post-exposure prophylaxis and curative treatment)

Ciprofloxacin may be used in the management of neutropenic patients with fever that is suspected to be due to a bacterial infection

Adults:

Broad spectrum antibiotic for infections caused by ciprofloxacin sensitive pathogens.

העלונים נשלחו לפרסום במאגר התרופות שבאתר האינטרנט של משרד הבריאות
וניתן לקבלו מודפס ע"י פניה לחברת טבע. <http://www.israeldrugs.health.gov.il>