



CIPRO TEVA 2 MG/ML - הודעה על הפסקת שיווק לצמיתות

תאריך: 17/12/2024 בעל הרישום טבע ישראל בע"מ מודיע על הפסקת שיווק לצמיתות של:

157 11 35047 00	מספר רישום
ציפרו-טבע 2 מ"ג/מ"ל	שם התכשיר בעברית
CIPRO-TEVA 2 MG/ML	שם התכשיר באנגלית
תמיסה לאנפוזיה	צורת מינון
תוך ורידי	דרך מתן
CIPROFLOXACIN AS LACTATE	מרכיב פעיל
2 MG / 1 ML	חוזק
Ciprofloxacin solution for infusion is indicated for the treatment of the following infections. Special attention should be paid to available information on resistance to ciprofloxacin before commencing therapy. Consideration should be given to official guidance on the appropriate use of antibacterial agents. Adults •Lower respiratory tract infections due to Gram-negative bacteria - exacerbations of chronic obstructive pulmonary disease. In exacerbation of chronic obstructive pulmonary disease Ciprofloxacin should be used only when it is considered inappropriate to use other antibacterial agents that are commonly recommended for the treatment of these infections. - broncho-pulmonary infections in cystic fibrosis or in bronchiectasis - pneumonia •Chronic suppurative otitis media •Acute exacerbation of chronic sinusitis especially if these are caused by Gram[1]negative bacteria •Urinary tract infections - Acute pyelonephritis - Complicated pyelonephritis - Bacterial prostatitis •□Genital tract infections - Epididymo-orchitis including cases due to susceptible Neisseria gonorrhoeae - Pelvic inflammatory disease including cases due to susceptible Neisseria gonorrhoeae •Infections of the gastro-intestinal tract (e.g. travellers' diarrhoea) •Intra-abdominal infections •Infections of the skin and soft tissue caused by Gram-negative bacteria •Malignant external otitis •Infections of the bones and joints •Inhalation anthrax (post-exposure prophylaxis and curative treatment)	התוויה

<p>Ciprofloxacin may be used in the management of neutropenic patients with fever that is suspected to be due to a bacterial infection</p> <p>Children and adolescents</p> <ul style="list-style-type: none"> •Broncho-pulmonary infections due to Pseudomonas aeruginosa in patients with cystic fibrosis •Complicated urinary tract infections and acute pyelonephritis •Inhalation anthrax (post-exposure prophylaxis and curative treatment) <p>Ciprofloxacin may also be used to treat severe infections in children and adolescents when this is considered to be necessary.</p> <p>Treatment should be initiated only by physicians who are experienced in the treatment of cystic fibrosis and/or severe infections in children and adolescents.</p>	
<p>אמצע יולי 2025</p>	<p>תאריך תחילת הפסקת השיווק</p>
<p>לצמיתות</p>	<p>תאריך צפי לסיום הפסקת השיווק</p>
	<p>תאריך חזרה לשיווק (יש למלא משבצת זאת עם חזרת התכשיר לשיווק בפועל)</p>
<p>לא</p>	<p>הכללה בסל הבריאות: כן / לא</p>
<p>מסחרית</p>	<p>סיבת הפסקת השיווק: סיבות תפעוליות / סיבות מסחריות</p>
<p>20 x 100 ML</p>	<p>גודל/י אריזה (נא לרשום את כל הגדלים הרלוונטיים להודעה זאת)</p>
	<p>זמינות של גודל אריזה אחר או חוזק אחר של התכשיר</p>

בברכה,

חגי קליין – רוקח ממונה

טבע ישראל בע"מ