

## SUMMARY OF PRODUCT CHARACTERISTICS

**TEVAGRASTIM® 30 MIU/0.5 ml**  
SOLUTION FOR INJECTION OR INFUSION  
IN PRE-FILLED SYRINGE

**TEVAGRASTIM® 48 MIU/0.8 ml**  
SOLUTION FOR INJECTION OR INFUSION  
IN PRE-FILLED SYRINGE

SUBCUTANEOUS INJECTION / INTRAVENOUS INFUSION

### 1. NAME OF THE MEDICINAL PRODUCT

Tevagrastim 30 MIU/0.5 mL solution for injection/infusion

Tevagrastim 48 MIU/0.8 mL solution for injection/infusion

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL of solution for injection/infusion contains 60 million international units [MIU] (600 µg) of filgrastim.

Each pre-filled syringe contains 30 MIU (300 µg) of filgrastim in 0.5 mL solution for injection/infusion.

Each pre-filled syringe contains 48 MIU (480 µg) of filgrastim in 0.8 mL solution for injection/infusion.

Filgrastim (recombinant methionyl human granulocyte colony stimulating factor) is produced in *Escherichia coli* K802 by recombinant DNA technology.

#### *Excipient with known effect*

Each mL of solution contains 50 mg of sorbitol.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Solution for injection/infusion

Clear, colourless solution.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Tevagrastim is indicated for the reduction in the duration of neutropenia and the incidence of febrile neutropenia in patients treated with established cytotoxic chemotherapy for malignancy (with the exception of chronic myeloid leukaemia and myelodysplastic syndromes) and for the reduction in the duration of neutropenia in patients undergoing myeloablative therapy followed by bone marrow transplantation considered to be at increased risk of prolonged severe neutropenia.

The safety and efficacy of filgrastim are similar in adults and children receiving cytotoxic chemotherapy.

Tevagrastim is indicated for the mobilisation of peripheral blood progenitor cells in normal donors (allogeneic PBPC).

In patients, children or adults, with severe congenital, cyclic, or idiopathic neutropenia with an absolute neutrophil count (ANC) of  $\leq 0.5 \times 10^9/L$ , and a history of severe or recurrent infections, long term administration of Tevagrastim is indicated to increase neutrophil counts and to reduce the incidence and duration of infection-related events.

Tevagrastim is indicated for the treatment of persistent neutropenia (ANC less than or equal to  $1.0 \times 10^9/L$ ) in patients with advanced HIV infection, in order to reduce the risk of bacterial infections when other options to manage neutropenia are inappropriate.

## **4.2 Posology and method of administration**

Filgrastim therapy should only be given in collaboration with an oncology centre which has experience in granulocyte colony stimulating factor (G-CSF) treatment and haematology and has the necessary diagnostic facilities. The mobilisation and apheresis procedures should be performed in collaboration with an oncology-haematology centre with acceptable experience in this field and where the monitoring of haematopoietic progenitor cells can be correctly performed.

### Established cytotoxic chemotherapy

#### *Posology*

The recommended dose of filgrastim is 0.5 MIU (5 µg)/kg/day. The first dose of filgrastim should be administered at least 24 hours after cytotoxic chemotherapy. . In randomised clinical trials, a subcutaneous dose of 23 MIU (230 µg)/m<sup>2</sup>/day (4.0 to 8.4 µg/kg/day) was used.

Daily dosing with filgrastim should continue until the expected neutrophil nadir is passed and the neutrophil count has recovered to the normal range. Following established chemotherapy for solid tumours, lymphomas and lymphoid leukaemias, it is expected that the duration of treatment required to fulfil these criteria will be up to 14 days. Following induction and consolidation treatment for acute myeloid leukaemia the duration of treatment may be substantially longer (up to 38 days) depending on the type, dose and schedule of cytotoxic chemotherapy used.

In patients receiving cytotoxic chemotherapy, a transient increase in neutrophil counts is typically seen 1 to 2 days after initiation of filgrastim therapy. However, for a sustained therapeutic response, filgrastim therapy should not be discontinued before the expected nadir has passed and the neutrophil count has recovered to the normal range. Premature discontinuation of filgrastim therapy prior to the time of the expected neutrophil nadir is not recommended.

#### *Method of administration*

Filgrastim may be given as a daily subcutaneous injection or as a daily intravenous infusion diluted in 5% glucose solution for infusion given over 30 minutes (see section 6.6). The subcutaneous route is preferred in most cases. There is some evidence from a study of single dose administration that intravenous dosing may shorten the duration of effect. The clinical relevance of this finding to multiple dose administration is not clear. The choice of route should depend on the individual clinical circumstance.

### In patients treated with myeloablative therapy followed by bone marrow transplantation

#### *Posology*

The recommended starting dose of filgrastim is 1.0 MIU (10 µg)/kg/day. The first dose of filgrastim should be administered at least 24 hours after cytotoxic chemotherapy and at least 24 hours after bone marrow infusion.

Once the neutrophil nadir has been passed the daily dose of filgrastim should be titrated against the neutrophil response as follows:

<b>Neutrophil count</b>	<b>Filgrastim dose adjustment</b>
> 1.0 x 10 <sup>9</sup> /L for 3 consecutive days	Reduce to 0.5 MIU (5 µg)/kg/day
Then, if ANC remains > 1.0 x 10 <sup>9</sup> /L for 3 more consecutive days	Discontinue filgrastim
If the ANC decreases to < 1.0 x 10 <sup>9</sup> /L during the treatment period the dose of filgrastim should be re-escalated according to the above steps	

*ANC = absolute neutrophil count*

#### *Method of administration*

Filgrastim may be given as a 30 minute or 24 hour intravenous infusion or given by continuous 24 hour subcutaneous infusion. Filgrastim should be diluted in 20 mL of 5% glucose solution for infusion (see section 6.6).

#### For the mobilisation of PBPCs in patients undergoing myelosuppressive or myeloablative therapy followed by autologous PBPC transplantation

##### *Posology*

The recommended dose of filgrastim for PBPC mobilisation when used alone is 1.0 MIU (10 µg)/kg/day for 5 to 7 consecutive days. Timing of leukapheresis: 1 or 2 leukapheresis on days 5 and 6 are often sufficient. In other circumstances, additional leukapheresis may be necessary. Filgrastim dosing should be maintained until the last leukapheresis.

The recommended dose of filgrastim for PBPC mobilisation after myelosuppressive chemotherapy is 0.5 MIU (5 µg)/kg/day from the first day after completion of chemotherapy until the expected neutrophil nadir is passed and the neutrophil count has recovered to the normal range. Leukapheresis should be performed during the period when the ANC rises from < 0.5 x 10<sup>9</sup>/L to > 5.0 x 10<sup>9</sup>/L. For patients who have not had extensive chemotherapy, one leukapheresis is often sufficient. In other circumstances, additional leukapheresis are recommended.

#### *Method of administration*

##### *Filgrastim for PBPC mobilisation when used alone:*

Filgrastim may be given as a 24 hour subcutaneous continuous infusion or subcutaneous injection. For infusions, filgrastim should be diluted in 20 mL of 5% glucose solution for infusion (see section 6.6).

##### *Filgrastim for PBPC mobilisation after myelosuppressive chemotherapy:*

Filgrastim should be given by subcutaneous injection.

#### For the mobilisation of PBPCs in normal donors prior to allogeneic PBPC transplantation

##### *Posology*

For PBPC mobilisation in normal donors, filgrastim should be administered at 1.0 MIU (10 µg)/kg/day for 4 to 5 consecutive days. Leukapheresis should be started at day 5 and continued until day 6 if needed in order to collect 4 x 10<sup>6</sup> CD34<sup>+</sup> cells/kg recipient bodyweight.

#### *Method of administration*

Filgrastim should be given by subcutaneous injection.

#### In patients with severe chronic neutropenia (SCN)

## *Posology*

### *Congenital neutropenia*

The recommended starting dose is 1.2 MIU (12 µg)/kg/day as a single dose or in divided doses.

### *Idiopathic or cyclic neutropenia*

The recommended starting dose is 0.5 MIU (5 µg)/kg/day as a single dose or in divided doses.

### *Dose adjustment*

Filgrastim should be administered daily by subcutaneous injection until the neutrophil count has reached and can be maintained at more than  $1.5 \times 10^9/L$ . When the response has been obtained, the minimal effective dose to maintain this level should be established. Long-term daily administration is required to maintain an adequate neutrophil count. After one to two weeks of therapy, the initial dose may be doubled or halved depending upon the patient's response. Subsequently the dose may be individually adjusted every 1 to 2 weeks to maintain the average neutrophil count between  $1.5 \times 10^9/L$  and  $10 \times 10^9/L$ . A faster schedule of dose escalation may be considered in patients presenting with severe infections. In clinical trials, 97 % of patients who responded had a complete response at doses of  $\leq 2.4$  MIU (24 µg)/kg/day. The long-term safety of filgrastim administration above 2.4 MIU (24 µg)/kg/day in patients with SCN has not been established.

### *Method of administration*

#### *Congenital, idiopathic or cyclic neutropenia:*

Filgrastim should be given by subcutaneous injection.

#### In patients with HIV infection

## *Posology*

### *For reversal of neutropenia*

The recommended starting dose of filgrastim is 0.1 MIU (1 µg)/kg/day with titration up to a maximum of 0.4 MIU (4 µg)/kg/day until a normal neutrophil count is reached and can be maintained (ANC  $> 2.0 \times 10^9/L$ ). In clinical studies,  $> 90$  % of patients responded at these doses, achieving reversal of neutropenia in a median of 2 days.

In a small number of patients ( $< 10$  %), doses up to 1.0 MIU (10 µg)/kg/day were required to achieve reversal of neutropenia.

### *For maintaining normal neutrophil counts*

When reversal of neutropenia has been achieved, the minimal effective dose to maintain a normal neutrophil count should be established. Initial dose adjustment to alternate day dosing with 30 MIU (300 µg)/day is recommended. Further dose adjustment may be necessary, as determined by the patient's ANC, to maintain the neutrophil count at  $> 2.0 \times 10^9/L$ . In clinical studies, dosing with 30 MIU (300 µg)/day on 1 to 7 days per week was required to maintain the ANC  $> 2.0 \times 10^9/L$ , with the median dose frequency being 3 days per week. Long-term administration may be required to maintain the ANC  $> 2.0 \times 10^9/L$ .

### *Method of administration*

Reversal of neutropenia or maintaining normal neutrophil counts: Filgrastim should be given by subcutaneous injection.

## Special populations

### *Elderly*

Clinical trials with filgrastim have included a small number of elderly patients but special studies have not been performed in this group and therefore specific dosage recommendations cannot be made.

### *Patients with renal or hepatic impairment*

Studies of filgrastim in patients with severe impairment of renal or hepatic function demonstrate that it exhibits a similar pharmacokinetic and pharmacodynamic profile to that seen in normal individuals. Dose adjustment is not required in these circumstances.

### *Paediatric use in the SCN and cancer settings*

Sixty-five percent of the patients studied in the SCN trial programme were under 18 years of age. The efficacy of treatment was clear for this age group, which included most patients with congenital neutropenia. There were no differences in the safety profiles for paediatric patients treated for SCN.

Data from clinical studies in paediatric patients indicate that the safety and efficacy of filgrastim are similar in both adults and children receiving cytotoxic chemotherapy.

The dosage recommendations in paediatric patients are the same as those in adults receiving myelosuppressive cytotoxic chemotherapy.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

Special warnings and precautions across indications

### *Hypersensitivity*

Hypersensitivity, including anaphylactic reactions, occurring on initial or subsequent treatment have been reported in patients treated with filgrastim. Permanently discontinue Tevagrastim in patients with clinically significant hypersensitivity. Do not administer filgrastim to patients with a history of hypersensitivity to filgrastim or pegfilgrastim.

### *Pulmonary adverse effects*

Pulmonary adverse reactions, in particular interstitial lung disease, have been reported after G-CSF administration. Patients with a recent history of lung infiltrates or pneumonia may be at higher risk. The onset of pulmonary signs such as cough, fever and dyspnoea in association with radiological signs of pulmonary infiltrates and deterioration in pulmonary function may be preliminary signs of acute respiratory distress syndrome (ARDS). Filgrastim should be discontinued and appropriate treatment given.

### *Glomerulonephritis*

Glomerulonephritis has been reported in patients receiving filgrastim and pegfilgrastim. Generally, events of glomerulonephritis resolved after dose reduction or withdrawal of filgrastim and pegfilgrastim. Urinalysis monitoring is recommended.

### *Capillary leak syndrome*

Capillary leak syndrome which can be life threatening if treatment is delayed, has been reported after G-CSF administration and is characterised by hypotension, hypoalbuminaemia, oedema and

haemoconcentration. Patients who develop symptoms of capillary leak syndrome should be closely monitored and receive standard symptomatic treatment, which may include a need for intensive care (see section 4.8).

#### *Splenomegaly and splenic rupture*

Generally asymptomatic cases of splenomegaly and cases of splenic rupture have been reported in patients and normal donors following administration of filgrastim. Some cases of splenic rupture were fatal. Therefore, spleen size should be carefully monitored (e.g. clinical examination, ultrasound). A diagnosis of splenic rupture should be considered in donors and/or patients reporting left upper abdominal or shoulder tip pain. Dose reductions of filgrastim have been noted to slow or stop the progression of splenic enlargement in patients with severe chronic neutropenia, and in 3% of patients a splenectomy was required.

#### *Malignant cell growth*

Granulocyte colony stimulating factor can promote growth of myeloid cells *in vitro* and similar effects may also be seen on some non myeloid cells *in vitro*.

#### *Myelodysplastic syndrome or chronic myeloid leukemia*

The safety and efficacy of filgrastim administration in patients with myelodysplastic syndrome or chronic myelogenous leukaemia have not been established. Filgrastim is not indicated for use in these conditions. Particular care should be taken to distinguish the diagnosis of blast transformation of chronic myeloid leukaemia from acute myeloid leukaemia.

#### *Acute myeloid leukaemia*

In view of limited safety and efficacy data in patients with secondary AML, filgrastim should be administered with caution.

The safety and efficacy of filgrastim administration in *de novo* AML patients aged < 55 years with good cytogenetics (t(8;21), t(15;17), and inv(16)) have not been established.

#### *Thrombocytopenia*

Thrombocytopenia has been reported in patients receiving filgrastim. Platelet counts should be monitored closely, especially during the first few weeks of filgrastim therapy. Consideration should be given to temporary discontinuation or dose reduction of filgrastim in patients with severe chronic neutropenia who develop thrombocytopenia (platelet count < 100 x 10<sup>9</sup>/l).

#### *Leukocytosis*

White blood cell counts of 100 x 10<sup>9</sup>/L or greater have been observed in less than 5 % of cancer patients receiving filgrastim at doses above 0.3 MIU/kg/day (3 µg/kg/day). No undesirable effects directly attributable to this degree of leukocytosis have been reported. However, in view of the potential risks associated with severe leukocytosis, a white blood cell count should be performed at regular intervals during filgrastim therapy. If leukocyte counts exceed 50 x 10<sup>9</sup>/L after the expected nadir, filgrastim should be discontinued immediately. However, during the period of administration of filgrastim for PBPC mobilisation, filgrastim should be discontinued or its dosage should be reduced if the leukocyte counts rise to > 70 x 10<sup>9</sup>/L.

#### *Immunogenicity*

As with all therapeutic proteins, there is a potential for immunogenicity. Rates of generation of antibodies against filgrastim is generally low. Binding antibodies do occur as expected with all biologics; however, they have not been associated with neutralising activity at present.

### *Aortitis*

Aortitis has been reported after G-CSF administration in healthy subjects and in cancer patients. The symptoms experienced included fever, abdominal pain, malaise, back pain and increased inflammatory markers (e.g. C reactive protein and white blood cell count). In most cases aortitis was diagnosed by CT scan and generally resolved after withdrawal of G-CSF. See also section 4.8.

### Special warnings and precautions associated with co-morbidities

#### *Special precautions in sickle cell trait and sickle cell disease*

Sickle cell crisis, in some cases fatal, have been reported with the use of filgrastim in patients with sickle cell trait or sickle cell disease. Physicians should use caution when prescribing filgrastim in patients with sickle cell trait or sickle cell disease.

#### *Osteoporosis*

Monitoring of bone density may be indicated in patients with underlying osteoporotic bone diseases who undergo continuous therapy with filgrastim for more than 6 months.

### Special precautions in cancer patients

*Filgrastim should not be used to increase the dose of cytotoxic chemotherapy beyond established dosage regimens.*

#### *Risks associated with increased doses of chemotherapy*

Special caution should be used when treating patients with high dose chemotherapy because improved tumour outcome has not been demonstrated and intensified doses of chemotherapeutic agents may lead to increased toxicities including cardiac, pulmonary, neurologic and dermatologic effects (please refer to the Summary of Product Characteristics of the specific chemotherapy agents used).

#### *Effect of chemotherapy on erythrocytes and thrombocytes*

Treatment with filgrastim alone does not preclude thrombocytopenia and anaemia due to myelosuppressive chemotherapy. Because of the potential of receiving higher doses of chemotherapy (e.g. full doses on the prescribed schedule) the patient may be at greater risk of thrombocytopenia and anaemia. Regular monitoring of platelet count and haematocrit is recommended. Special care should be taken when administering single or combination chemotherapeutic agents which are known to cause severe thrombocytopenia.

The use of filgrastim mobilised PBPCs has been shown to reduce the depth and duration of thrombocytopenia following myelosuppressive or myeloablative chemotherapy.

#### *Myelodysplastic syndrome and acute myeloid leukaemia in breast and lung cancer patients*

In the post-marketing observational study setting, myelodysplastic syndrome (MDS) and acute myeloid leukaemia (AML) have been associated with the use of pegfilgrastim, an alternative G-CSF medicine, in conjunction with chemotherapy and/or radiotherapy in breast and lung cancer patients. A similar association between filgrastim and MDS/AML has not been observed. Nonetheless, patients with breast cancer and patients with lung cancer should be monitored for signs and symptoms of MDS/AML.

### *Other special precautions*

The effects of filgrastim in patients with substantially reduced myeloid progenitors have not been studied. Filgrastim acts primarily on neutrophil precursors to exert its effect in elevating neutrophil counts. Therefore, in patients with reduced precursors neutrophil response may be diminished (such as those treated with extensive radiotherapy or chemotherapy, or those with bone marrow infiltration by tumour).

Vascular disorders, including veno-occlusive disease and fluid volume disturbances, have been reported occasionally in patients undergoing high dose chemotherapy followed by transplantation.

There have been reports of graft versus host disease (GvHD) and fatalities in patients receiving G-CSF after allogeneic bone marrow transplantation (see sections 4.8 and 5.1).

Increased haematopoietic activity of the bone marrow in response to growth factor therapy has been associated with transient abnormal bone scans. This should be considered when interpreting bone-imaging results.

### Special precautions in patients undergoing PBPC mobilisation

#### *Mobilisation*

There are no prospectively randomised comparisons of the two recommended mobilisation methods (filgrastim alone or in combination with myelosuppressive chemotherapy) within the same patient population. The degree of variation between individual patients and between laboratory assays of CD34<sup>+</sup> cells mean that direct comparison between different studies is difficult. It is therefore difficult to recommend an optimum method. The choice of mobilisation method should be considered in relation to the overall objectives of treatment for an individual patient.

#### *Prior exposure to cytotoxic agents*

Patients who have undergone very extensive prior myelosuppressive therapy may not show sufficient mobilisation of PBPC to achieve the recommended minimum yield ( $\geq 2.0 \times 10^6$  CD34<sup>+</sup> cells/kg) or acceleration of platelet recovery to the same degree.

Some cytotoxic agents exhibit particular toxicities to the haematopoietic progenitor pool and may adversely affect progenitor mobilisation. Agents such as melphalan, carmustine (BCNU) and carboplatin, when administered over prolonged periods prior to attempts at progenitor mobilisation, may reduce progenitor yield. However, the administration of melphalan, carboplatin or BCNU together with filgrastim has been shown to be effective for progenitor mobilisation. When a PBPC transplantation is envisaged it is advisable to plan the stem cell mobilisation procedure early in the treatment course of the patient. Particular attention should be paid to the number of progenitors mobilised in such patients before the administration of high-dose chemotherapy. If yields are inadequate, as measured by the criteria above, alternative forms of treatment not requiring progenitor support should be considered.

#### *Assessment of progenitor cell yields*

In assessing the number of progenitor cells harvested in patients treated with filgrastim, particular attention should be paid to the method of quantitation. The results of flow cytometric analysis of CD34<sup>+</sup> cell numbers vary depending on the precise methodology used and recommendations of numbers based on studies in other laboratories need to be interpreted with caution.

Statistical analysis of the relationship between the number of CD34<sup>+</sup> cells re-infused and the rate of platelet recovery after high-dose chemotherapy indicates a complex but continuous relationship.



The recommendation of a minimum yield of  $\geq 2.0 \times 10^6$  CD34<sup>+</sup> cells/kg is based on published experience resulting in adequate haematologic reconstitution. Yields in excess of this appear to correlate with more rapid recovery, those below with slower recovery.

#### Special precautions in normal donors undergoing PBPC mobilisation

Mobilisation of PBPC does not provide a direct clinical benefit to normal donors and should only be considered for the purposes of allogeneic stem cell transplantation.

PBPC mobilisation should be considered only in donors who meet normal clinical and laboratory eligibility criteria for stem cell donation with special attention to haematological values and infectious disease.

The safety and efficacy of filgrastim have not been assessed in normal donors < 16 years or > 60 years.

Transient thrombocytopenia (platelets <  $100 \times 10^9/L$ ) following filgrastim administration and leukapheresis was observed in 35 % of subjects studied. Among these, two cases of platelets <  $50 \times 10^9/L$  were reported and attributed to the leukapheresis procedure.

If more than one leukapheresis is required, particular attention should be paid to donors with platelets <  $100 \times 10^9/L$  prior to leukapheresis; in general apheresis should not be performed if platelets <  $75 \times 10^9/L$ .

Leukapheresis should not be performed in donors who are anticoagulated or who have known defects in haemostasis.

Donors who receive G-CSFs for PBPC mobilisation should be monitored until haematological indices return to normal.

#### Special precautions in recipients of allogeneic PBPCs mobilised with filgrastim

Current data indicate that immunological interactions between the allogeneic PBPC graft and the recipient may be associated with an increased risk of acute and chronic GvHD when compared with bone marrow transplantation.

#### Special precautions in SCN patients

Filgrastim should not be administered to patients with severe congenital neutropenia who develop leukaemia or have evidence of leukaemic evolution.

#### *Blood cell counts*

Other blood cell changes occur, including anaemia and transient increases in myeloid progenitors, which require close monitoring of cell counts

#### *Transformation to leukaemia or myelodysplastic syndrome*

Special care should be taken in the diagnosis of SCNs to distinguish them from other haematopoietic disorders such as aplastic anaemia, myelodysplasia and myeloid leukaemia. Complete blood cell counts with differential and platelet counts and an evaluation of bone marrow morphology and karyotype should be performed prior to treatment.

There was a low frequency (approximately 3 %) of myelodysplastic syndromes (MDS) or leukaemia in clinical trial patients with SCN treated with filgrastim. This observation has only been made in patients with congenital neutropenia. MDS and leukaemias are natural complications of the disease and are of uncertain relation to filgrastim therapy. A subset of approximately 12 % of patients who had

normal cytogenetic evaluations at baseline was subsequently found to have abnormalities, including monosomy 7, on routine repeat evaluation. It is currently unclear whether long-term treatment of patients with SCN will predispose patients to cytogenetic abnormalities, MDS or leukaemic transformation. It is recommended to perform morphologic and cytogenetic bone marrow examinations in patients at regular intervals (approximately every 12 months).

#### *Other special precautions*

Causes of transient neutropenia such as viral infections should be excluded.

Haematuria was common and proteinuria occurred in a small number of patients. Regular urinalysis should be performed to monitor these events.

The safety and efficacy in neonates and patients with autoimmune neutropenia have not been established.

#### Special precautions in patients with HIV infection

##### *Blood cell counts*

ANC should be monitored closely, especially during the first few weeks of filgrastim therapy. Some patients may respond very rapidly and with a considerable increase in neutrophil count to the initial dose of filgrastim. It is recommended that the ANC is measured daily for the first 2 to 3 days of filgrastim administration. Thereafter, it is recommended that the ANC is measured at least twice weekly for the first two weeks and subsequently once per week or once every other week during maintenance therapy. During intermittent dosing with 30 MIU (300 µg)/day of filgrastim, there can be wide fluctuations in the patient's ANC over time. In order to determine a patient's trough or nadir ANC, it is recommended that blood samples are taken for ANC measurement immediately prior to any scheduled dosing with filgrastim.

##### *Risk associated with increased doses of myelosuppressive medicinal products*

Treatment with filgrastim alone does not preclude thrombocytopenia and anaemia due to myelosuppressive medicinal products. As a result of the potential to receive higher doses or a greater number of these medicinal products with filgrastim therapy, the patient may be at higher risk of developing thrombocytopenia and anaemia. Regular monitoring of blood counts is recommended (see above).

##### *Infections and malignancies causing myelosuppression*

Neutropenia may be due to bone marrow infiltrating opportunistic infections such as *Mycobacterium avium* complex or malignancies such as lymphoma. In patients with known bone marrow-infiltrating infections or malignancy, consider appropriate therapy for treatment of the underlying condition in addition to administration of filgrastim for treatment of neutropenia. The effects of filgrastim on neutropenia due to bone marrow-infiltrating infection or malignancy have not been well established.

#### All patients

##### *Sorbitol*

Tevagrastim contains sorbitol (E420). The additive effect of concomitantly given products containing sorbitol (or fructose) and dietary intake of sorbitol (or fructose) should be taken into account.

##### *For intravenous administration*

Patients with hereditary fructose intolerance (HFI) must not be given this medicinal product unless strictly necessary. Infants and young children (below 2 years of age) may not yet be diagnosed with HFI. Medicinal products (containing sorbitol/fructose) given intravenously may be life-threatening

and should not be given in this population unless there is an overwhelming clinical need and no alternatives are available.

A detailed history of HFI symptoms has to be taken of each patient before giving this medicinal product.

#### *Sodium*

This medicinal product contains less than 1 mmol sodium (23 mg) per pre-filled syringe, i.e. essentially 'sodium-free'.

#### *Traceability*

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

### **4.5 Interaction with other medicinal products and other forms of interaction**

The safety and efficacy of filgrastim given on the same day as myelosuppressive cytotoxic chemotherapy have not been definitively established. In view of the sensitivity of rapidly dividing myeloid cells to myelosuppressive cytotoxic chemotherapy, the use of filgrastim is not recommended in the period from 24 hours before to 24 hours after chemotherapy. Preliminary evidence from a small number of patients treated concomitantly with filgrastim and 5-Fluorouracil indicates that the severity of neutropenia may be exacerbated.

Possible interactions with other haematopoietic growth factors and cytokines have not yet been investigated in clinical trials.

Since lithium promotes the release of neutrophils, it is likely to potentiate the effect of filgrastim. Although this interaction has not been formally investigated, there is no evidence that such an interaction is harmful.

### **4.6 Fertility, pregnancy and lactation**

#### *Pregnancy*

There are no or limited amount of data from the use of filgrastim in pregnant women. Studies in animals have shown reproductive toxicity. An increased incidence of embryo-loss has been observed in rabbits at high multiples of the clinical exposure and in the presence of maternal toxicity (see section 5.3). There are reports in the literature where the transplacental passage of filgrastim in pregnant women has been demonstrated.

Filgrastim is not recommended during pregnancy.

#### *Breast-feeding*

It is unknown whether filgrastim/metabolites are excreted in human milk. A risk to the breastfed newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from filgrastim therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

#### *Fertility*

Filgrastim did not affect reproductive performance or fertility in male or female rats (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Filgrastim may have a minor influence on the ability to drive and use machines. Dizziness may occur following the administration of filgrastim (see section 4.8).

## **4.8 Undesirable effects**

### **a. Summary of the safety profile**

The most serious adverse reactions that may occur during filgrastim treatment include: anaphylactic reaction, serious pulmonary adverse events (including interstitial pneumonia and ARDS), capillary leak syndrome, severe splenomegaly/splenic rupture, transformation to myelodysplastic syndrome or leukaemia in SCN patients, GvHD in patients receiving allogeneic bone marrow transfer or peripheral blood cell progenitor cell transplant and sickle cell crisis in patients with sickle cell disease.

The most commonly reported adverse reactions are pyrexia, musculoskeletal pain (which includes bone pain, back pain, arthralgia, myalgia, pain in extremity, musculoskeletal pain, musculoskeletal chest pain, neck pain), anaemia, vomiting, and nausea. In clinical trials in cancer patients musculoskeletal pain was mild or moderate in 10%, and severe in 3% of patients.

### **b. Tabulated summary of adverse reactions**

The data in the table below describe adverse reactions reported from clinical trials and spontaneous reporting.

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

MedDRA system organ class	Adverse reactions			
	Very common (≥ 1/10)	Common (≥ 1/100 to < 1/10)	Uncommon (≥ 1/1,000 to < 1/100)	Rare (≥ 1/10,000 to < 1/1,000)
Infections and Infestations		Sepsis Bronchitis Upper respiratory tract infection Urinary tract infection		
<i>Blood and lymphatic system disorders</i>	Thrombocytopenia Anaemia	Splenomegaly <sup>a</sup> Haemoglobin decreased <sup>c</sup>	Leukocytosis <sup>a</sup>	Splenic rupture <sup>a</sup> Sickle cell anaemia with crisis Extramedullary haematopoiesis
<i>Immune system disorders</i>			Hypersensitivity Drug Hypersensitivity <sup>a</sup> Graft versus Host Disease <sup>b</sup>	Anaphylactic reaction
<i>Metabolism and nutrition disorders</i>		Decreased appetite <sup>c</sup> Blood lactate dehydrogenase increased	Hyperuricaemia Blood uric acid increased	Blood glucose decreased Pseudogout <sup>a</sup> (Chondrocalcinosis Pyrophosphate) Fluid volume disturbances
Psychiatric disorders		Insomnia		
<i>Nervous system disorders</i>	Headache <sup>a</sup>	Dizziness Hypoaesthesia Paraesthesia		
<i>Vascular disorders</i>		Hypertension Hypotension	Veno-occlusive disease <sup>d</sup>	Capillary leak syndrome <sup>a</sup> Aortitis
<i>Respiratory, thoracic and mediastinal disorders</i>		Haemoptysis Dyspnoea Cough <sup>a</sup> Oropharyngeal pain <sup>a,e</sup> Epistaxis	Acute respiratory distress syndrome <sup>a</sup> Respiratory failure <sup>a</sup> Pulmonary oedema <sup>a</sup> Pulmonary haemorrhage Interstitial lung disease <sup>a</sup> Lung infiltration <sup>a</sup> Hypoxia	
<i>Gastro-intestinal disorders</i>	Diarrhoea <sup>a, c</sup> Vomiting <sup>a, e</sup> Nausea <sup>a</sup>	Oral pain Constipation <sup>c</sup>		

<i>Hepato-biliary disorders</i>		Hepatomegaly Blood alkaline phosphatase increased	Aspartate Amino-transferase increased Gamma-glutamyl transferase increased	
<i>Skin and sub-cutaneous tissue disorders</i>	Alopecia <sup>a</sup>	Rash <sup>a</sup> Erythema	Rash maculopapular	Cutaneous vasculitis <sup>a</sup> Sweets syndrome (acute febrile neutrophilic dermatosis)
<i>Musculo-skeletal and connective tissue disorders</i>	Musculoskeletal pain <sup>c</sup>	Muscle spasms	Osteoporosis	Bone density decreased Exacerbation of rheumatoid arthritis
<i>Renal and urinary disorders</i>		Dysuria Haematuria	Proteinuria	Glomerulonephritis Urine abnormality
<i>General disorders and administration site conditions</i>	Fatigue <sup>a</sup> Mucosal inflammation <sup>a</sup> Pyrexia	Chest pain <sup>a</sup> Pain <sup>a</sup> Asthenia <sup>a</sup> Malaise <sup>c</sup> Oedema peripheral <sup>c</sup>	Injection site reaction	
<i>Injury, poisoning and procedural complications</i>		Transfusion reaction <sup>c</sup>		

<sup>a</sup> See section c (Description of selected adverse reactions)

<sup>b</sup> There have been reports of GvHD and fatalities in patients after allogeneic bone marrow transplantation (see section c)

<sup>c</sup> Includes bone pain, back pain, arthralgia, myalgia, pain in extremity, musculoskeletal pain, musculoskeletal chest pain, neck pain

<sup>d</sup> Cases were observed in the post-marketing setting in patients undergoing bone marrow transplant or PBPC mobilisation

<sup>e</sup> Adverse events with higher incidence in filgrastim patients compared to placebo and associated with the sequelae of the underlying malignancy or cytotoxic chemotherapy

### c. Description of selected adverse reactions

#### *Hypersensitivity*

Hypersensitivity-type reactions including anaphylaxis, rash, urticaria, angioedema, dyspnoea and hypotension occurring on initial or subsequent treatment have been reported in clinical studies and in post marketing experience. Overall, reports were more common after IV administration. In some cases, symptoms have recurred with rechallenge, suggesting a causal relationship. Tevagrastim should be permanently discontinued in patients who experience a serious allergic reaction.

#### *Pulmonary adverse events*

In clinical studies and the post-marketing setting pulmonary adverse effects including interstitial lung

disease, pulmonary oedema, and lung infiltration have been reported in some cases with an outcome of respiratory failure or acute respiratory distress syndrome (ARDS), which may be fatal (see section 4.4).

#### *Splenomegaly and splenic rupture*

Cases of splenomegaly and splenic rupture have been reported following administration of filgrastim. Some cases of splenic rupture were fatal (see section 4.4).

#### *Capillary leak syndrome*

Cases of capillary leak syndrome have been reported with granulocyte colony-stimulating factor use. These have generally occurred in patients with advanced malignant diseases, sepsis, taking multiple chemotherapy medications or undergoing apheresis (see section 4.4).

#### *Cutaneous vasculitis*

Cutaneous vasculitis has been reported in patients treated with filgrastim. The mechanism of vasculitis in patients receiving filgrastim is unknown. During long term use cutaneous vasculitis has been reported in 2% of SCN patients.

#### *Leukocytosis*

Leukocytosis ( $WBC > 50 \times 10^9/l$ ) was observed in 41% of normal donors and transient thrombocytopenia (platelets  $< 100 \times 10^9/l$ ) following filgrastim and leukapheresis was observed in 35% of donors (see section 4.4).

#### *Sweets syndrome*

Cases of Sweets syndrome (acute febrile neutrophilic dermatosis) have been reported in patients treated with filgrastim.

#### *Pseudogout (chondrocalcinosis pyrophosphate)*

Pseudogout (chondrocalcinosis pyrophosphate) has been reported in patients with cancer treated with filgrastim.

#### *GvHD*

There have been reports of GvHD and fatalities in patients receiving G-CSF after allogeneic bone marrow transplantation (see sections 4.4 and 5.1).

#### d. Paediatric population

Data from clinical studies in paediatric patients indicate that the safety and efficacy of filgrastim are similar in both adults and children receiving cytotoxic chemotherapy suggesting no age-related differences in the pharmacokinetics of filgrastim. The only consistently reported adverse event was musculoskeletal pain, which is no different from the experience in the adult population.

There is insufficient data to further evaluate filgrastim use in paediatric subjects.

#### e. Other special populations

##### *Geriatric use*

No overall differences in safety or effectiveness were observed between subjects over 65 years of age compared to younger adult (>18 years of age) subjects receiving cytotoxic chemotherapy and clinical experience has not identified differences in the responses between elderly and younger adult patients.

There is insufficient data to evaluate filgrastim use in geriatric subjects for other approved filgrastim indications.

##### *Paediatric SCN patients*

Cases of decreased bone density and osteoporosis have been reported in paediatric patients with SCN receiving chronic treatment with filgrastim.

##### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il/>

#### **4.9 Overdose**

The effects of filgrastim overdosage have not been established.

Discontinuation of filgrastim therapy usually results in a 50 % decrease in circulating neutrophils within 1 to 2 days, with a return to normal levels in 1 to 7 days.



## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Immunostimulants, colony stimulating factors, ATC code: L03AA02

Tevagrastim is a biosimilar medicinal product.

Human G-CSF is a glycoprotein which regulates the production and release of functional neutrophils from the bone marrow. Tevagrastim containing r-metHuG-CSF (filgrastim) causes marked increases in peripheral blood neutrophil counts within 24 hours, with minor increases in monocytes. In some SCN patients, filgrastim can also induce a minor increase in the number of circulating eosinophils and basophils relative to baseline; some of these patients may present with eosinophilia or basophilia prior to treatment. Elevations of neutrophil counts are dose-dependent at recommended doses. Neutrophils produced in response to filgrastim show normal or enhanced function as demonstrated by tests of chemotactic and phagocytic function. Following termination of filgrastim therapy, circulating neutrophil counts decrease by 50 % within 1 to 2 days, and to normal levels within 1 to 7 days.

Use of filgrastim in patients undergoing cytotoxic chemotherapy leads to significant reductions in the incidence, severity and duration of neutropenia and febrile neutropenia. Treatment with filgrastim significantly reduces the duration of febrile neutropenia, antibiotic use and hospitalisation after induction chemotherapy for acute myelogenous leukaemia or myeloablative therapy followed by bone marrow transplantation. The incidence of fever and documented infections were not reduced in either setting. The duration of fever was not reduced in patients undergoing myeloablative therapy followed by bone marrow transplantation.

Use of filgrastim, either alone or after chemotherapy, mobilises haematopoietic progenitor cells into peripheral blood. These autologous PBPCs may be harvested and infused after high-dose cytotoxic therapy, either in place of or in addition to bone marrow transplantation. Infusion of PBPCs accelerates haematopoietic recovery reducing the duration of risk for haemorrhagic complications and the need for platelet transfusions.

Recipients of allogeneic PBPCs mobilised with filgrastim experienced significantly more rapid haematological recovery, leading to a significant decrease in time to unsupported platelet recovery, when compared with allogeneic bone marrow transplantation.

One retrospective European study evaluating the use of G-CSF after allogeneic bone marrow transplantation in patients with acute leukaemias suggested an increase in the risk of GvHD, treatment related mortality (TRM) and mortality when G-CSF was administered. In a separate retrospective international study in patients with acute and chronic myelogenous leukaemias, no effect on the risk of GvHD, TRM and mortality was seen. A meta-analysis of allogeneic transplant studies, including the results of nine prospective randomised trials, 8 retrospective studies and 1 case-controlled study, did not detect an effect on the risks of acute GvHD, chronic GvHD or early treatment-related mortality.

<b>Relative risk (95 % CI) of GvHD and TRM following treatment with G-CSF after bone marrow transplantation</b>					
<i>Publication</i>	<i>Period of study</i>	<i>N</i>	<i>Acute grade II-IV GvHD</i>	<i>Chronic GvHD</i>	<i>TRM</i>
Meta-analysis (2003)	1986-2001 <sup>a</sup>	1198	1.08 (0.87, 1.33)	1.02 (0.82, 1.26)	0.70 (0.38, 1.31)
European retrospective study (2004)	1992-2002 <sup>b</sup>	1789	1.33 (1.08, 1.64)	1.29 (1.02, 1.61)	1.73 (1.30, 2.32)
International retrospective study (2006)	1995-2000 <sup>b</sup>	2110	1.11 (0.86, 1.42)	1.10 (0.86, 1.39)	1.26 (0.95, 1.67)
<sup>a</sup> Analysis includes studies involving bone marrow transplant during this period; some studies used GM-CSF (granulocyte-macrophage colony stimulating factor)					
<sup>b</sup> Analysis includes patients receiving bone marrow transplant during this period					

Prior to allogeneic PBPC transplantation, use of filgrastim for the mobilisation of PBPC in normal donors allows a collection of  $\geq 4 \times 10^6$  CD34<sup>+</sup> cells/kg recipient body weight in the majority of the donors after two leukapheresis. Normal donors are given a dose of a 10 µg/kg/day, administered subcutaneously for 4 to 5 consecutive days.

Use of filgrastim in patients, children or adults, with SCN (severe congenital, cyclic, and idiopathic neutropenia) induces a sustained increase in absolute neutrophil counts in peripheral blood and a reduction of infection and related events.

Use of filgrastim in patients with HIV infection maintains normal neutrophil counts to allow scheduled dosing of antiviral and/or other myelosuppressive medication. There is no evidence that patients with HIV infection treated with filgrastim show an increase in HIV replication.

As with other haematopoietic growth factors, G-CSF has shown *in vitro* stimulating properties on human endothelial cells.

The efficacy and safety of Tevagrastim has been assessed in randomised, controlled phase III studies in breast cancer, lung cancer and Non-Hodgkin-Lymphoma. There were no relevant differences between Tevagrastim and the reference product with regard to duration of severe neutropenia and incidence of febrile neutropenia.

## **5.2 Pharmacokinetic properties**

Randomised, single-blind, single dose, crossover studies in 196 healthy volunteers showed that the pharmacokinetic profile of Tevagrastim was comparable to that of the reference product after subcutaneous and intravenous administration.

Clearance of filgrastim has been shown to follow first-order pharmacokinetics after both subcutaneous and intravenous administration. The serum elimination half-life of filgrastim is approximately 3.5 hours, with a clearance rate of approximately 0.6 mL/min/kg. Continuous infusion with filgrastim over a period of up to 28 days, in patients recovering from autologous bone marrow transplantation, resulted in no evidence of drug accumulation and comparable elimination half-lives. There is a positive linear correlation between the dose and the serum concentration of filgrastim, whether administered intravenously or subcutaneously. Following subcutaneous administration of recommended doses, serum concentrations were maintained above 10 ng/mL for 8 to 16 hours. The volume of distribution in blood is approximately 150 mL/kg.

In cancer patients, the pharmacokinetic profile of Tevagrastim and the reference product was comparable after single and repeated subcutaneous administration.

### **5.3 Preclinical safety data**

Filgrastim was studied in repeated dose toxicity studies up to 1 year in duration which revealed changes attributable to the expected pharmacological actions including increases in leukocytes, myeloid hyperplasia in bone marrow, extramedullary granulopoiesis and splenic enlargement. These changes all reversed after discontinuation of treatment.

Effects of filgrastim on pre-natal development have been studied in rats and rabbits. Intravenous (80 µg/kg/day) administration of filgrastim to rabbits during the period of organogenesis was maternally toxic and increased spontaneous abortion, post-implantation loss, and decreased mean live litter size and foetal weight were observed.

Based on reported data for another filgrastim product similar to the reference product, comparable findings plus increased foetal malformations were observed at 100 µg/kg/day, a maternally toxic dose which corresponded to a systemic exposure of approximately 50-90 times the exposures observed in patients treated with the clinical dose of 5 µg/kg/day. The no observed adverse effect level for embryo-foetal toxicity in this study was 10 µg/kg/day, which corresponded to a systemic exposure of approximately 3-5 times the exposures observed in patients treated with the clinical dose.

In pregnant rats, no maternal or foetal toxicity was observed at doses up to 575 µg/kg/day. Offspring of rats administered filgrastim during the peri-natal and lactation periods, exhibited a delay in external differentiation and growth retardation ( $\geq 20$  µg/kg/day) and slightly reduced survival rate (100 µg/kg/day).

Filgrastim had no observed effect on the fertility of male or female rats.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sorbitol (E420)  
Glacial acetic acid  
Polysorbate 80  
Sodium hydroxide (for pH-adjustment)  
Water for injections

### **6.2 Incompatibilities**

Tevagrastim must not be diluted with sodium chloride solution.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

Diluted filgrastim may be adsorbed to glass and plastic materials except diluted, as described in section 6.6.

### **6.3 Shelf life**

The expiry date of the product is indicated on the packaging materials.

After dilution: Chemical and physical in-use stability of the diluted solution for infusion has been demonstrated for 24 hours at 2 °C to 8 °C. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 °C to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

## 6.4 Special precautions for storage

Store in a refrigerator (2 °C – 8 °C).

Patient can store the medicine out of refrigerator below 25°C, for up to 5 days, without exceeding the expiry date. Once removed from the refrigerator, the medicine should be used within 5 days or be disposed of. The medicine should not be returned to the refrigerator.

For storage conditions after dilution of the medicinal product, see section 6.3.

## 6.5 Nature and contents of container

Pre-filled syringe (type I glass) with injection needle (stainless steel), with or without a needle safety guard.

### Tevagrastim 30 MIU/0.5 ml solution for injection or infusion.

Packs containing 1, 5 or 10 pre-filled syringes with a safety device, with 0.5 ml solution for injection or infusion; or multipacks containing 10 (2 packs of 5) pre-filled syringes with a safety device, with 0.5 ml solution for injection or infusion.

### Tevagrastim 48 MIU/0.8 ml solution for injection or infusion.

Packs containing 1, 5 or 10 pre-filled syringes with a safety device, with 0.8 ml solution for injection or infusion; or multipacks containing 10 (2 packs of 5) pre-filled syringes with a safety device, with 0.8 ml solution for injection or infusion.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling

If required, Tevagrastim may be diluted in glucose 50 mg/mL (5 %) solution for infusion.

Dilution to a final concentration less than 0.2 MIU (2 µg) per mL is not recommended at any time.

The solution should be visually inspected prior to use. Only clear solutions without particles should be used.

For patients treated with filgrastim diluted to concentrations below 1.5 MIU (15 µg) per mL, human serum albumin (HSA) should be added to a final concentration of 2 mg/mL.

Example: In a final injection volume of 20 mL, total doses of filgrastim less than 30 MIU (300 µg) should be given with 0.2 mL of 200 mg/mL (20 %) human albumin solution added.

When diluted in glucose 50 mg/mL (5 %) solution for infusion, Tevagrastim is compatible with glass and a variety of plastics including PVC, polyolefin (a co-polymer of polypropylene and polyethylene) and polypropylene.

Tevagrastim does not contain any preservative. In view of the possible risk of microbial contamination, Tevagrastim syringes are for single use only.

Accidental exposure to freezing temperatures does not adversely affect the stability of Tevagrastim.

### Using the pre-filled syringe with a needle safety guard

The needle safety guard covers the needle after injection to prevent needle stick injury. This does not affect normal operation of the syringe. Depress the plunger slowly and evenly until the entire dose has been given and the plunger cannot be depressed any further. While maintaining pressure on the plunger, remove the syringe from the patient. The needle safety guard will cover the needle when releasing the plunger.

## Disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. LICENCE HOLDER AND MANUFACTURER**

Teva Israel Ltd.  
124 Dvora HaNevi'a St., Tel Aviv 6944020 Israel

### **8. REGISTRATION NUMBER**

143.29.31990

Revised in October 2024.